

EINSTEIN CLUSTER ALIGNMENTS REVISITED

SCOTT W. CHAMBERS¹, ADRIAN L. MELOTT¹, AND CHRISTOPHER J. MILLER²

Draft version February 1, 2008

ABSTRACT

We have examined whether the major axes of rich galaxy clusters tend to point toward their nearest neighboring cluster. We have used the data of Ulmer, McMillan, and Kowalski, who used position angles based on X-ray morphology. We also studied a subset of this sample with updated positions and distances from the MX Northern Abell Cluster Survey (for rich clusters ($R \geq 1$) with well known redshifts). A Kolmogorov-Smirnov (KS) test showed no significant signal for nonrandom angles on any scale $\leq 100h^{-1}\text{Mpc}$. However, refining the null hypothesis with the Wilcoxon rank-sum test, we found a high confidence signal for alignment. Confidence levels increase to a high of 99.997% as only near neighbors which are very close are considered. We conclude there is a strong alignment signal in the data, consistent with gravitational instability acting on Gaussian perturbations.

Subject headings: galaxies: clusters: general — large-scale structure of universe

1. INTRODUCTION

It is well documented that clusters of galaxies tend to be elongated, elliptical systems (eg., Carter & Metcalfe 1980) giving them major axes and “position angles” in the sky. Bingelli (1982) found that the major axes of rich galaxy clusters have the tendency (in projection) to point toward their nearest neighbor cluster whose distance, d_n , was closer than $\sim 30h^{-1}\text{Mpc}$. Since then, there have been multiple studies on whether this “Bingelli effect” actually exists. Much of this literature supports the reality of the effect. Flin (1987) and Rhee & Katgert (1987) both found significant alignments for d_n less than $\sim 30h^{-1}\text{Mpc}$. West (1989) and Rhee, van Haaren & Katgert (1992) have also detected a signal for alignment. Plionis (1994) found weak alignment signals up to $d_n \sim 60h^{-1}\text{Mpc}$, with more significant cluster alignments on smaller scales ($10\text{--}30h^{-1}\text{Mpc}$). Moreover, West, Jones and Forman (1995) found evidence that galaxy cluster substructure tends to be aligned with its host cluster and surrounding environment out to $\sim 10h^{-1}\text{Mpc}$, which might help explain these alignments.

Not all authors favor an alignment effect, however. Both Struble & Peebles (1985) and Ulmer, McMillan & Kowalski (1989) (hereafter UMK) found no significant evidence that clusters point toward their nearest neighbor (see however, Argyles et al. 1986).

Galactic positions may not be good tracers of the shape of a cluster, for galaxies contribute discreteness noise. Most clusters contain much more mass in hot, X-ray emitting gas than that of the galaxies themselves. Dark matter contributes more mass to the system than gas and galaxies combined. Thus, the shape of the actual cluster mass cannot be directly seen. However, it is believed that the X-ray emitting gas within a cluster traces its gravitational potential (Sarazin 1986). X-ray morphology is, then, probably the best observable for determining galaxy cluster shape and orientation.

Cluster alignments are not crucial in distinguishing cos-

mological models, but they are additional evidence in support of the gravitational instability hypothesis of structure formation (Shandarin & Klypin 1984; Splinter *et al.* 1997; Onuora & Thomas 2000).

For these reasons, the negative results of UMK are interesting. Whereas most authors used galaxies to define ellipticity and spatial orientation, both UMK and West *et al.* chose to use X-ray morphology. UMK did not find a statistical alignment for any nearest neighbor distance scale. There are many more papers (including West *et al.*) that found an alignment effect than didn’t. Since UMK used the shape of the X-ray gas in their search for alignment, their negative results are even more important. Most analyses of numerical simulations of structure formation by gravitational instability from Gaussian initial perturbations predict alignments on some scale, which provides some physical motivation for detecting the alignments searched for by UMK.

Both Onuora & Thomas 2000 and Splinter *et al.* 1997, for example, predicted alignments for d_n of at least $15h^{-1}\text{Mpc}$ for standard CDM models. Onuora & Thomas extended this distance to $30h^{-1}\text{Mpc}$ for ΛCDM models. Both these studies showed that the predicted alignment differences for individual cosmological background models are not practical means for determining cosmological parameters, such as Ω . However, Splinter *et al.* presented evidence that cluster ellipticity and the scale dependence of cluster alignments probe the primordial power spectrum independent of the parameters of the background cosmology. The alignments in these simulations fit a general picture of cluster formation by hierarchical clustering in which material falls into the cluster along the large scale filamentary structure, as interpreted by Shandarin & Klypin (1984). This picture has been supported by dynamical evidence of drainage along such filaments (Novikov *et al.* 1999).

2. DATA AND ANALYSIS

¹Dept. of Physics & Astronomy, Univ. of Kansas, Lawrence, KS 66045

²Dept. of Physics & Astronomy, Univ. of Maine, Orono, ME 04469

2.1. Subject Clusters and Position Angles

UMK determined the major axis orientation of 46 X-ray clusters observed by *Einstein* (UMK Table 1). The major axis served to define a position angle on the celestial sphere (measured counter-clock-wise from north). UMK used both $R = 0$ and $R \geq 1$ Abell clusters in their original analysis. However, $R = 0$ clusters were never part of Abell’s (1958) statistical sample and should not be used in nearest-neighbor analyses (since one needs to be sure of the existence of both the source cluster and its nearest neighbor). UMK created smaller subsets of clusters (out of their original 46) for analysis as well. We will re-examine only the largest (46 cluster) subset from UMK, since this was the only one which was large enough to use after we eliminated clusters for which we did not have good redshift information. We analyzed the data in UMK Table 1 as given, again with updated redshifts (results we show here), and then re-analyzed a subset of it which met our stringent selection criteria as described below.

2.2. Potential Neighbor Sample

We start out with the 46 clusters from UMK Table 1. However, over the past ten years many clusters have new or revised redshifts, and so we also searched an updated Abell cluster redshift survey to revise any of the nearest neighbors in UMK and also to apply constraints that will limit any selection effects and biases in our data. Our angles are measured counter-clockwise from North as in UMK. Our distances are measured for a Friedmann Universe with $q_0 = 0$ and $h = H_0/100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

The recent success of galaxy cluster surveys (Miller *et al.* 1999; 2000 and references therein) gives us a large and uniform sample, as compared with previous efforts. The cluster redshifts and coordinates we have used are mainly from the MX Northern Abell Cluster Survey (MX) (Slinglend *et al.* 1998; Miller *et al.* 2000). There are 256 Abell $R \geq 1$ clusters (i.e. $\text{dec.} \geq -27^\circ$) having measured redshifts and within $0.012 \leq z \leq 0.10$ and $|b| \geq 30^\circ$. These clusters have an average of 25 measured galaxies each, and 90% have more than one measured redshift. Having multiple redshifts per cluster is quite important, as the elongation in redshift space due to internal velocity dispersion in the cluster is easily comparable to typical nearest neighbor distances. Miller *et al.* (1999) found that cluster redshifts based on only one galaxy redshift are erroneous by $\pm 500 \text{ km s}^{-1}$ 41% of the time. While this error may seem small ($\sim 5h^{-1} \text{ Mpc}$), such an error can easily throw off the determination of the nearest-neighbor in dense environments. This northern hemisphere sample is nearly complete to $z = 0.10$ (e.g. Miller & Batuski 2000). To define potential nearest neighbors, we searched through these 256 clusters noting the distance and direction to the nearest neighbor and the distance to the nearest edge of the survey. We then made a potentially important cut: If the boundary of the survey region were closer than the nearest neighbor, the pair would not be used for our alignment statistics. This is because there is a potential nearer neighbor hidden outside the boundary.

After we applied the above constraints to the original UMK data, we were left with 25 clusters out of their original 46. We call this our Statistical Sample and we present our data in Table 1. We also found that ten of these 25

clusters now have *different* nearest-neighbors than those found by UMK. In four of these ten cases, UMK used an $R = 0$ cluster as their nearest neighbor. In other words, the original UMK dataset remains relatively unchanged, with 40/46 clusters having the same nearest-neighbor in our new analysis. It is worth noting that the constraints that we apply to the original UMK data prevent a large number of biases from entering the analysis. For example, by using only $R \geq 1$ clusters we are ensuring that our base cluster subset is a statistically complete sample. We also use clusters with multiple-galaxy determined redshifts so that we can be sure of their location. Finally, perhaps our most important improvement over the UMK data, is that we exclude clusters where the edge of the survey is closer than the nearest-neighbor.

2.3. Analysis

We first analyze the UMK data as given (in their Table 1) with revised nearest neighbors for six clusters. Our purpose in re-analyzing the original UMK data is to determine whether their statistical analyses were sensitive enough to actually detect a cluster alignment. We note that the UMK dataset contains $R = 0$ clusters and does not exclude clusters that have a survey-edge which is closer than the nearest neighbor. We then analyze our subset of 25 clusters which meet our stringent selection criteria designed to produce a uniform well-controlled sample.

UMK have provided a position angle for the major axis of the cluster as measured via the X-ray emission. As this is an orientation, not a direction, the angle between it and the projected direction to the nearest neighbor, the pointing angle ϕ_p can only have a range $-90 \leq \phi_p \leq 90$. We also assumed that the sign is not significant, so we examine $0 \leq |\phi_p| \leq 90$. This coincides with the UMK procedure.

We therefore define alignment as a tendency for the angles $|\phi_p|$ to be *smaller* than they would be if distributed isotropically, that is, uniformly over this interval. Most previous work has not really tested for alignment - rather, it has tested for *any* kind of anisotropy.

2.3.1. Re-analysis of UMK Data

We repeated the Kolmogorov-Smirnov (KS) test as used by UMK to test against a distribution sampled from a population uniformly distributed over this interval, as ϕ_p would be if there are no correlations. The KS test (Lehman 1975) is used to test against the null hypothesis that the sample (our angles ϕ_p) could be drawn from a parent population of ϕ_p . We have no *a priori* reason to believe that the parent distribution of ϕ_p ’s should be anything other than random. In other words, the null hypothesis for the KS test is that our observed sample is the same as our parent (random) distribution. The KS test measures the significance of the null hypothesis being false. Although widely used in the “cluster alignment” literature, it in fact is a test for non-uniformity (since our hypothetical parent population has a uniform distribution of ϕ_p). The KS test uses the maximum value of the difference between cumulative distribution functions as its diagnostic. Thus, KS is excellent at detecting any deviation of the sample from the parent population. However, this very attribute makes it weaker at detecting a more specific signal (in our case the signal of alignment) than some other tests. Using the KS

test, our results agree with UMK. We find that the null hypothesis can be discounted with only 73% confidence. This is too small a confidence level to rule out the null hypothesis. We have also looked only at those clusters with d_n below some critical value, d_c . For d_c down to $10h^{-1}\text{Mpc}$, the confidence for ruling out the null hypothesis is still too small.

Refinement of the null hypothesis makes an enormous difference. We are not looking for just *any* difference, we are looking for alignments, which means the angles ϕ_p are systematically lower than they would be if drawn from a uniform parent population of ϕ . The Wilcoxon rank-sum test (WRS) (Lehmann 1975) tests for this. The null hypothesis of WRS is that the sample is not systematically smaller or larger than the parent population. Thus, while WRS is more sensitive to alignment (small angles), it would not (for example) be sensitive to a tendency for the pointing angles to clump around 45 degrees. The WRS ranks the populations and is thus sensitive to alignment differences between the real and assumed parent population. The WRS returns a signed result, indicating that our sample has systematically higher or lower ϕ_p 's. By defining more precisely what is tested, a great increase in statistical power can be achieved.

Using WRS on the updated UMK data, we found no significant effect. However, if we use it only on the UMK clusters with $d_n \leq 30h^{-1}\text{Mpc}$, we find (for the 38 UMK clusters that meet this criterion) 99.8% confidence in alignment. Restriction to smaller distances produces alignments with much higher confidence, as seen in Table 2. Refinement of the null hypothesis combined with restrictions to nearer neighbors produces a strong signal. We also note that the original UMK data (i.e. with six erroneous nearest neighbors) also shows a similarly strong significance (99.6%) for alignment. In other words, had UMK used a more targeted statistical tool (such as the WRS), they too would have certainly noted this alignment effect.

2.3.2. Analysis of Revised UMK Data

We next examined the set of our 25 clusters with corrected distances in our Statistical Sample. We observe that this set has fractionally fewer nearest neighbors at large distances than in UMK's Table 1, suggesting corrections of misidentification of near neighbors in that. However, we still found no significant signal with the KS test. Applying WRS to our entire controlled sample of 25 clusters produced, as it did with the entire UMK sample, no substantial confidence in alignment. Restricting the study to the clusters with $d_n \leq 20h^{-1}\text{Mpc}$ leaves 17 clusters, with a confidence level of 98.4%. More restrictive limits on distance produces increased confidence, up to very high levels, as with the UMK data.

Table 2 shows the results of the WRS test applied to

both the original UMK data (with six revised cluster neighbors) as well as to the 25 clusters with corrected distances and stringent constraints to account for any biasing or selection effects. Table 2 shows the strength of the alignment effect as we vary the restrictions on how near the neighbors must be in order to be considered.

There are suggestive trends in the data. Although the UMK data contain nearly twice as many clusters, as more restrictive distance cuts are applied, the number of surviving clusters converge until both sets are nearly the same size for $d_n \leq 10h^{-1}\text{Mpc}$. Because clusters have correlated spatial positions, a random error in cluster position measurement is much more likely to move a cluster away from a near neighbor than to create a spurious one. Our more well-controlled sample has a smaller mean nearest neighbor distance.

The UMK result, in this case, nevertheless shows a strong signal when even moderately restrictive cuts are made. By making these cuts we are most likely removing erroneous near neighbor identifications. The surviving close near neighbors in UMK may well be correct, in spite of being at low galactic latitude (where obscuration is a problem), being in poorly sampled regions, or being too close to a survey boundary.

3. CONCLUSIONS AND DISCUSSION

Re-examining the UMK *Einstein* cluster data, we find that testing for alignment (with WRS) rather than for *any* departure from uniformity of angles (as with KS) allows us to find a significant signal for alignment in this data for clusters with nearest neighbors at distances $d_n \leq 30h^{-1}\text{Mpc}$. When we use a more stringently defined sample, we still find a strong signal for alignment, reaching 3.34σ when we restrict $d_n \leq 10h^{-1}\text{Mpc}$. Thus, refinement of the null hypothesis has proven crucial in finding the alignment signal. Use of the well-controlled sample was in this case not necessary to find a signal, but it confirms (with a lower confidence due to a smaller sample size) that the X-ray emission from galaxy clusters does tend to point to the nearest cluster neighbor.

A potential weakness in most alignment studies is the search for nearest neighbor alignment rather than supercluster axis alignment. In the future, we plan to examine a larger sample using the supercluster axis finding procedure defined in Novikov et al. (1999). The work we present here serves to remove an inconsistency from the data analysis and help clarify some issues of statistical methodology.

ALM wishes to acknowledge support from the National Science Foundation under grant number AST-0070702, the University of Kansas General Research Fund and the National Center for Supercomputing Applications.

REFERENCES

- Abell, G. O. 1958, ApJS, 3, 211
- Argyres, P. C., Groth, E. J., Peebles, P. J. E., Struble, M. F. 1986, ApJ, 91, 471
- Binggelli, B. 1982, A&A, 107, 338
- Carter, D., and Metcalfe, N. 1979, MNRAS, 191, 325
- Flin P. 1987, MNRAS, 228, 941
- Lehmann, E.L. 1975 Nonparametrics: Statistical Methods Based on Ranks (Prentice Hall: New Jersey)
- Miller, C., & Batuski, D. 2000, submitted to ApJ (astro-ph/0002295)
- Miller, C.J., Batuski, D.J., Slingland, K.A., and Hill, J.M. 1999 ApJ 523, 492
- Miller, C.J., Krughoff, K.S., Batuski, D.J., Slingland, K.A., & Hill, J.M. 2000, submitted to AJ
- Novikov, D.I., Melott, A.L., Wilhite, B.B., Kaufman, M., Burns, J.O., Miller, C.J., and Batuski, D.J., 1999 MNRAS, 304, L5
- Onuora, L.I. and Thomas, P.A., 2000 MNRAS in press (astro-ph/0003402)
- Plionis, M., 1994, ApJ 95, 401

TABLE 1
DATA SUMMARY FOR STATISTICAL SAMPLE

Cluster					Nearest Neighbor					
Abell no.	R.A.	dec.	Dist.	ϕ	Abell no.	R.A.	dec.	Dist.	d_n	ϕ_p
85	0.652	-9.617	162	166	87 ^a	0.675	-10.067	161	2.1	23.6
119	0.897	-1.533	130	18	168	1.210	-0.017	132	11.5	54.1
154	1.138	17.400	185	57	150	1.110	12.900	172	19.6	51.7
168	1.210	-0.017	132	156	119	0.897	-1.533	130	11.5	83.9
399	2.920	12.817	210	31	401	2.937	13.383	214	4.3	7.8
401	2.937	13.383	214	38	399	2.920	12.817	210	4.3	13.8
1367	11.698	20.117	65	137	1656	12.957	28.250	69	22.3	70.3
1656	12.957	28.250	69	49	1367	22.330	11.698	65	22.3	17.7
1767	13.570	59.467	204	145	1904	14.338	48.783	205	44.9	12.2
1775	13.660	26.617	208	117	1795 ^b	13.778	26.833	184	24.8	34.0
1795	13.778	26.833	184	21	1831 ^b	13.948	28.233	179	9.7	40.2
1809	13.847	5.400	228	1	1780 ^a	13.702	3.133	227	12.4	42.8
1904	14.338	48.783	205	84	1767	13.570	59.467	204	44.9	48.8
1991	14.870	18.833	171	56	1913	14.408	16.900	154	25.7	18.4
2029	15.142	5.950	224	131	2028	15.118	7.717	225	7.1	37.5
2063	15.343	8.817	104	55	2147	16.000	16.033	103	21.7	1.2
2065	15.343	27.900	210	151	2056 ^a	15.285	28.450	216	7.0	28.7
2079	15.433	29.050	192	41	2092	15.522	31.317	194	8.9	10.5
2107	15.627	21.933	121	167	2152 ^b	16.052	16.583	121	17.0	37.0
2124	15.718	36.217	192	135	2122 ^a	15.710	36.283	192	2.7	16.2
2147	16.000	16.033	103	159	2151 ^a	16.050	17.883	108	5.9	43.1
2151	16.050	17.883	108	116	2147 ^a	16.000	16.033	103	5.9	86.1
2152	16.052	16.583	121	113	2151	16.050	17.883	108	13.0	65.7
2199	16.448	39.633	88	43	2197	16.442	41.017	91	3.4	46.7
2670	23.860	-10.683	221	104	2659 ^a	23.708	-15.750	226	22.0	79.8

NOTE.—Key to columns is as follows:
Col. (1). – Abell cluster number.
Cols. (2)-(3). – Right ascension (hours) and declination (degrees) of the cluster (1950 epoch).
Col. (4). – Distance to the cluster (h^{-1} Mpc).
Col. (5). – Position angle of the cluster.
Cols. (6) - (10) – For the nearest neighbor cluster.
Col. (6). – Number of the Abell cluster.
Cols. (7) - (8). – Right ascension (hours) and declination (degrees) of the cluster (1950 epoch).
Col. (9). – Distance to the cluster (h^{-1} Mpc).
Col. (10). – Distance between the cluster (1) and its nearest neighbor (6) (h^{-1} Mpc).
Col. (11). – “Pointing” angle in degrees; see text for definition.
^a Different nearest neighbor than that found by UMK.
^b Different nearest neighbor than that found by UMK. However, UMK used an $R = 0$ cluster.

TABLE 2
WILCOXON RANK-SUM RESULTS

UMK Data (rev.)			Stat. Sample	
D_{max} (h^{-1} Mpc)	Sig.	N_{cl}	Sig.	N_{cl}
10	4.00 σ (99.997%)	12	3.34 σ (99.96%)	11
20	3.45 σ (99.97%)	29	2.14 σ (98.4%)	17
30	2.90 σ (99.8%)	38	1.27 σ (89.8%)	23
None	0.94 σ (82.6%)	46	0.87 σ (80.8%)	25




FIG. 1.— The distribution of position angle against nearest neighbor distance is shown for the UMK original 46 clusters. Six of these clusters have revised nearest neighbor distances and pointing angles.




FIG. 2.— The distribution of position angle against nearest neighbor distance is shown for our sample of 25 clusters, using UMK position angles with corrected distances than were found in UMK.

Rhee G.F.R.N., Katgert P., 1987, A&A, 183, 217
 Rhee G.F.R.N., van Haarlem M., Katgert P., 1992, AJ, 103, 1721
 Sarazin, C.L., 1986, Rev. Mod. Phys., 58, 1
 Shandarin, S.F., and Klypin, A.A. 1984, Sov Astron. 28, 491
 Slingend, K., Batuski, D., Miller, C., Haase, S., Michaud, K., & Hill, J. 1998, ApJS, 115, 1

Splinter R.J., Melott A.L., Linn A.M., Buck C., Tinker J., 1997, ApJ 479, 632
 Struble M.F., Peebles P.J.E., 1985, AJ, 90, 592
 Ulmer M.P., McMillan S.L.W., Kowalski M.P., 1989, ApJ, 338, 711
 West M.J., 1989, ApJ, 347, 610
 West M.J., Jones C., Forman W., 1995, ApJ, 451, L5